

Name (PRINT) KEY

(35 points total)

1. If the same amount of heat is added to 5.00 g samples of the metals listed below, which metal will experience the **largest temperature change**? [3 pts]

Metal	Specific Heat Capacity (J/g·°C)
Al	0.897
Au	0.129
Cu	0.385
Fe	0.449
K	0.753

- a) Al                      b) **Au \***                      c) Cu                      d) Fe                      e) K
2. How much **heat** needs to be added to a 4.33 g piece of aluminum to raise its temperature from 22.0°C to 27.0°C? [ $s_{\text{Al}} = 0.900 \text{ J/g}\cdot\text{°C}$ ] [3 pts]
- a) 19.5 J \*                      b) 21.7 J                      c) 24.1 J                      d) 0.0416 J                      e) none of these

$$q = ms\Delta T$$

$$q = (4.33 \text{ g})(0.900 \text{ J/g}\cdot\text{°C})(27.0\text{°C} - 22.0\text{°C}) = \mathbf{19.5 \text{ J}}$$

3. A piece of stainless steel ( $s_{\text{steel}} = 0.50 \text{ J/g}\cdot\text{°C}$ ) is transferred from an oven (183°C) to 125 g of water at 23.2°C, into which it is immersed. The water and steel together reach a temperature of 51.5°C. What is the **mass** of the steel? [ $s_{\text{H}_2\text{O}} = 4.184 \text{ J/g}\cdot\text{°C}$ ] [4 pts]

$$-q_{\text{ss}} = +q_{\text{water}}$$

$$-m_{\text{ss}}s_{\text{ss}}\Delta T_{\text{ss}} = +q_{\text{w}}s_{\text{w}}\Delta T_{\text{w}}$$

$$-(m_{\text{ss}})(0.50 \text{ J/g}\cdot\text{°C})(51.5\text{°C} - 183\text{°C}) = +(125 \text{ g})(4.184 \text{ J/g}\cdot\text{°C})(51.5\text{°C} - 23.2\text{°C})$$

$$65.75m_{\text{ss}} = 1.48 \times 10^4$$

$$m_{\text{ss}} = \mathbf{225 \text{ g}}$$

4. A particular form of electromagnetic radiation has a frequency of  $6.32 \times 10^{14} \text{ 1/s}$ .
- a) What is its **wavelength** ( $\lambda$ ) in **nanometers**? [3 pts]

$$c = \lambda\nu$$

$$3.00 \times 10^8 \text{ m/s} = \lambda(6.32 \times 10^{14} \text{ 1/s})$$

$$\lambda = 4.75 \times 10^{-7} \text{ m} \times \frac{1 \text{ nm}}{1 \times 10^{-9} \text{ m}} = \mathbf{4.75 \times 10^2 \text{ nm} = 475 \text{ nm}}$$

- b) What is the **energy** (in joules) of *one photon* of this radiation? [2 pts]

$$E = h\nu$$

$$E = (6.626 \times 10^{-34} \text{ J}\cdot\text{s})(6.32 \times 10^{14} \text{ 1/s}) = \mathbf{4.19 \times 10^{-19} \text{ J}}$$

- c) What is the **energy** of a *mole* of photons with the above frequency? [2 pts]

$$\frac{4.19 \times 10^{-19} \text{ J}}{1 \text{ photon}} \times \frac{6.022 \times 10^{23} \text{ photons}}{1 \text{ mol}} = \mathbf{2.52 \times 10^5 \text{ J/mol}}$$

5. **TRUE or FALSE**

a) Blue light has a longer wavelength than red light. [2 pts]

**FALSE**, blue light is higher energy than red light and hence has a shorter wavelength.

b) The energy of electromagnetic radiation is *directly* proportional to the frequency of the radiation. [2 pts]

**TRUE**

6. If the frequency of electromagnetic radiation **decreases**, which of the following statements is **correct**? [3 pts]

a) The number of cycles passing a given point per unit time increases.

b) The velocity of the radiation decreases.

c) The wavelength of the radiation decreases.

d) The energy of the radiation increases.

e) **None of the above is correct.** \*

7. An electron in an excited state of the hydrogen atom, drops from the  $n = 7$  energy state to the  $n = 4$  energy state. Which of the following statements is **true**? [3 pts]

a) This change is called absorption.

b) The *wavelength* of light emitted when an electron moves from the  $n = 7$  to the  $n = 4$  energy state is **shorter** than the wavelength emitted when an electron moves from the  $n = 7$  to the  $n = 1$  energy state.

c) **The magnitude of the energy change in going from  $n = 7$  to  $n = 4$  is the same as going from  $n = 4$  to  $n = 7$ .**

d) The  $n = 7$  to  $n = 4$  electronic transition in a hydrogen atom will have the same energy change as the  $n = 7$  to  $n = 4$  transition in a helium atom.

e) None of the above is true.

8. The energy (in joules) of an electron energy level in the Bohr atom is given by the expression:

$$E_n = -2.18 \times 10^{-18} \text{ J} \left( \frac{1}{n^2} \right) \text{ where } n \text{ is the principal quantum number for the energy level.}$$

a) Calculate the **energy** (in J) of both the 1<sup>st</sup> energy level and the 6<sup>th</sup> energy level in a hydrogen atom? [3 pts]

$$E_1 = -2.18 \times 10^{-18} \text{ J} \left( \frac{1}{1^2} \right) = -2.18 \times 10^{-18} \text{ J}$$

$$E_6 = -2.18 \times 10^{-18} \text{ J} \left( \frac{1}{6^2} \right) = -6.06 \times 10^{-20} \text{ J}$$

b) Calculate  $\Delta E$  between the 1<sup>st</sup> and 6<sup>th</sup> energy levels in the hydrogen atom. [2 pts]

$$\Delta E = E_f - E_i$$

$$\Delta E = -6.06 \times 10^{-20} \text{ J} - (-2.18 \times 10^{-18} \text{ J}) = 2.12 \times 10^{-18} \text{ J}$$

c) What is the **wavelength** of the photon necessary to promote an electron from  $n = 1$  to  $n = 6$  in a hydrogen atom? [3 pts]

$$E = \frac{hc}{\lambda}$$

$$\lambda = \frac{hc}{E} = \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})(3.00 \times 10^8 \text{ m/s})}{2.12 \times 10^{-18} \text{ J}} = 9.38 \times 10^{-8} \text{ m}$$

**Potentially Useful Information**

$$q = ms\Delta T$$

$$\Delta T = T_f - T_i$$

$$s_{\text{H}_2\text{O}} = 4.184 \text{ J/g}\cdot^\circ\text{C}$$

$$c = \lambda\nu$$

$$E = h\nu$$

$$E = \frac{hc}{\lambda}$$

$$c = 3.00 \times 10^8 \text{ m/s}$$

$$h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$$

$$E_n = -R_H \left( \frac{1}{n^2} \right)$$

$$R_H = 2.18 \times 10^{-18} \text{ J}$$

$$\Delta E = E_f - E_i$$